maintaining the data needed, and c including suggestions for reducing	lection of information is estimated to ompleting and reviewing the collect this burden, to Washington Headqu uld be aware that notwithstanding an DMB control number.	ion of information. Send comments arters Services, Directorate for Info	s regarding this burden estimate ormation Operations and Reports	or any other aspect of the s, 1215 Jefferson Davis	his collection of information, Highway, Suite 1204, Arlington
1. REPORT DATE 30 SEP 1997		2. REPORT TYPE		3. DATES COVERED <b>00-00-1997 to 00-00-1997</b>	
4. TITLE AND SUBTITLE				5a. CONTRACT NUMBER	
Lagrangian Floats for Deep Convection				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)  University of Washington, Applied Physics Laboratory, 1013 N.E. 40th  Street, Seattle, WA, 98105				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAII Approved for publ	ABILITY STATEMENT ic release; distributi	on unlimited			
13. SUPPLEMENTARY NO	OTES				
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFIC		17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON	
a. REPORT unclassified	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE unclassified	Same as Report (SAR)	2	

**Report Documentation Page** 

Form Approved OMB No. 0704-0188

### LAGRANGIAN FLOATS FOR DEEP CONVECTION

Eric A. D'Asaro
Applied Physics Laboratory
University of Washington
1013 NE 40th Str
Seattle, WA 98105
dasaro@apl.washington.edu
206 685 2982
206 543 6785 (fax)
Award No. N000149410025

# LONG TERM GOALS

I aim to understand the process of deep convection in the ocean.

#### **OBJECTIVES**

Near surface water is mixed to great depth at a few high latitude locations, thereby forming the deep and bottom masses of the ocean. This proposal supports the development and deployment of neutrally buoyant floats which will accurately follow the three-dimensional motion of water parcels in these regions of Deep Convection, and the deployment of these floats in the Labrador Sea in the winters 1997 and 1998. During deep convection, the floats move vertically with the convective motions, thus directly measuring where, when, how and how fast the surface water is converted to deep water.

### **APPROACH**

The floats to be used are a variant of existing instruments, but designed for longer duration and deeper depths. They accurately follow water motions through a combination of a density which matches that of seawater and a high drag. The density is matched to that of the ambient water by actively changing the float's volume and will stay matched, despite changes in pressure and temperature, though a combination of active control and a hull compressibility which is close to that of seawater. High drag is achieved through a large circular drogue attached to the float. The horizontal motion of the float is determined by acoustic tracking (RAFOS) and its vertical motion is determined from pressure. Data is relayed at the end of the 2 month mission via satellite (ARGOS).


# **ACCOMPLISHMENTS**

13 floats were deployed in February, 1997 in the Labrador Sea. Of these, 11 returned full data records. 3 floats were recovered, 2 by fishermen, and returned to APL for diagnosis. Overall, the floats worked very well. Several technical problems were found and we expect even better performance in the winter 1998 deployments.

### SCIENTIFIC/TECHNICAL RESULTS

The Figure 1 shows depth/time data from all of the floats. A convectively mixed layer extending from the surface to about 700m is evident. For 20 days the floats cycle the floats cycle across this layer, with speeds often exceeding 10 cm/s. The depth of the convecting layer appears to be set by the intrusion of Irminger Sea water at this depth.

Occasionally, the floats descend through this barrier, reaching depths of about 1 km. Two



excursions to 1000 m occur. The rms vertical velocity of 2.3 cm/s is approximately that expected for convection with typical heat fluxes. Temperature measurements on the floats clearly show a vertical heat flux comparable to that expected.

Overall, the data clearly show strong mixing to 700m and intermittant mixing to 1 km. The mixing time, equal to the time for a float to traverse the layer, is about a day.

### IMPACT FOR SCIENCE

These direct measurement of the vertical velocity and temperature fluctuations associated with deep convection will allow direct comparisons with theory and numerical models. The successful development of long-lived autonomous Lagrangian floats should allow similar measurements in a variety of strongly mixing regimes in both the Littoral and open ocean.

### **RELATED PROJECTS**

These floats are close relatives of those used to study mixing in the Littoral zone and upper ocean mixed layer funded by ONR 322PO. Mixing processes in these various environments are similar in many ways and we learn the most by comparing and contrasting them.